

Corrigendum — *Telopea* 11 (3)

Veronica sect. *Labiatooides* Wettst. not ‘*V. sect. Derwentia*’

Briggs BG & Ehrendorfer F (2006) New Australian species and typifications in *Veronica* sens. lat. (Plantaginaceae). *Telopea* 11 (3) 276–292.

Briggs BG & Ehrendorfer F (2006) Chromosome numbers of Australian and New Guinean species of *Veronica* (Plantaginaceae). *Telopea* 11 (3) 294–298.

In these papers we discussed the evolution of *Veronica* in Australia, using a proposed sectional name (‘*V. sect. Derwentia*’) for the clade of *Veronica* that includes most of the Australian members. This sectional name was not formally published. Since these papers went to press we have become aware that there is, however, an existing name applicable to this group, i.e. *Veronica* sect. *Labiatooides* Wettst. A type was not designated by Wettstein (1891, page 86) but lectotypification is being made in Garnock-Jones et al. (2007).

References

- Garnock-Jones P, Albach D & Briggs B (2007) Botanical names in Southern Hemisphere *Veronica* (Plantaginaceae) sect. *Detzneria*, sect. *Hebe*, and sect. *Labiatooides*. *Taxon* 56 (in press).
- Wettstein R (1891) Scrophulariaceae. In Engler HGA & Prantl KAE, *Die Natürlichen Pflanzenfamilien* 4 (3B) 39–107.

Barbara G. Briggs¹ & F. Ehrendorfer²

¹*Botanic Gardens Trust Sydney, Mrs Macquaries Road, Sydney NSW 2000, Australia.*

²*Institute of Botany, University of Vienna, Rennweg 14, A-1030, Vienna, Austria.*